



1. Name of the supplement.
2. Supplement form.
3. Usage.
4. How to use SLAMR™?
5. Special precautions & Warnings.
6. Ingredients.
7. Nutritional value.
8. Influence of ability to drive and use machinery.
9. Interactions with medicine or other interactions.
10. Durability.
11. Storage conditions.
12. Packaging (type and content).
13. Side effects.
14. Contact.

What is SLAMR™?

SLAMR™ is a product developed with the purpose of boosting sexual performance for men. It is based on a natural formula from herbs and plants. The idea behind SLAMR™ is to remove the limitations you set for yourself, by doing this the product inspires the user to accomplish more and help the user unleash his full potential.

1. Name of the supplement:

SLAMR™

2. Supplement form:

Dark brown gel/jelly.

3. Usage:

SLAMR™ is improving sexual performance for men.

- Increases the chance for a harder and longer lasting erection.
- Increases your ability for repeated sexual activity.
- Enables a longer, more intense orgasm.
- Increases excitement and energy during intercourse.
- Reduces the risk of premature ejaculations.
- Increases sex drive.

4. How to use SLAMR™?

- Take one packet (12 grams) and consume it orally, swallow it.
- You can take it with water or other liquids, but it is not necessary.
- Take it 30 minutes before sexual intercourse.
- Recommended temperature of the product should be over 20 degrees celsius to easily consume it and to get it out of the packet.
- *Do not exceed the limit of 1 packet or 12 grams per 24 hours.*

5. Special precautions & Warnings:

- Do not exceed the limit of 1 packet or 12 grams per 24 hours.
- Do not use if you have a heart condition.
- Do not use if you suffer from high or low blood pressure.
- Consult your healthcare practitioner for advice and direction before use.
- Keep out of reach of children.
- Not for use by persons under the age of 18.

People should not take SLAMR™ if they have:

- A hormone-sensitive cancer or condition.
- A known sensitivity to Epimedium Brevicornum.
- Are taking aromatase inhibitors.
- Been recommended no to do so by a doctor.
- A Bleeding disorder.
- A Surgery: *can increase the risk of bleeding during surgery.*
- Diabetes: *talk to your doctor before using SLAMR™.*
- Seizures: *If you have ever had a seizure, don't use SLAMR™.*
- Mental conditions: *Use with caution.*
- Kidney problems.
- Diarrhea.
- Glaucoma.
- Asthma or allergies: *if you have bee allergy or pollen allergy, don't use SLAMR™.*
- Liver disease.

6. Ingredients:

Glucose Syrup(78,61%), Straine Flower Honey(11,12%), Mulberry Molasses(3,78%), Bee Pollen(1,57%), Epimedium Brevicornum(0,79%), Carob(0,63%), Oat(0,63%), Ginger(0,63%), Lesser Galangal(0,47%), Cinnamon(0,31%), Maca(0,31%), American Ginseng(0,16%), Nettle(0,16%), Siberian Ginseng(0,13%), Ginkgo Biloba(0,13%), Red Ginseng(0,13%), Pumpkin(0,11%), Tribulus Terrestris(0,11%), Kola Nut(0,08%), Vanilla Flavor(0,08%), Royal Jelly(0,06%).

7. Nutritional value:

Nutritional value per packet (12 Grams)	
Recommended daily dosage is 12 Grams.	
Calories	42,4 kcal
Total fat	76,8 mg
Saturated fat	37,2 mg
Trans fat	<6 mg
Protein	276 mg
Total carbohydrates	10,4 g
Sugar	9,4 g
Dietary fiber	447,6 mg
Sodium	9,6 mg

8. Influence of ability to drive and use machinery:

SLAMR™ should not affect your ability to drive or operate machinery.

9. Interactions with medicine or other interactions.

Any blood thinner or blood pressure medicine should not be taken with SLAMR™.

Any heart medications or other medications for erectile dysfunction should not be taken with SLAMR™.

If you are using any medications regularly consult with your medical practitioner before using SLAMR™.

List of medications or substances that may interact:

Ace Inhibitors, Adenosine, Alprazolam,
Aluminum Hydroxide, Amitriptyline, Amlodipine, Amphetamines, Anastrozole,
Anesthesia, Antiarrhythmics, Antibiotics, Antidepressants, Antihistamines,
Aromatase Inhibitors, Aspirin.
Beta-blockers, Bumetanide, Buspirone.
Calcium Carbonate, Calcium Channel Blockers, Captopril, Carbamazepine,
Carisoprodol, Celecoxib, Cetirizine, Chlorpropamide, Cimetidine, Ciprofloxacin,
Citalopram, Clarithromycin, Clonazepam, Clopidogrel, Clozapine, Codeine,
Cortisone, Cyclobenzaprine, Cyclosporine, Cytochrome P450 Substrates.
Dalteparin, Decongestants, Desipramine, Dexamethasone, Diazepam,
Diclofenac, Digoxin, Dihydroxy Aluminum. Sodium Carbonate, Diltiazem,
Diphenhydramine, Dipyridamole, Disulfiram, Diuretics, Donepezil.
Efavirenz, Enalapril, Enoxacin, Enoxaparin, Ephedrine, Esomeprazole,
Estradiol, Estrogens, Ethinyl Estradiol, Exemestane.
Famotidine, Felodipine, Fentanyl, Fexofenadine, Flecainide, Fluconazole,
Fluoxetine, Fluoxetine, Fluvastatin, Fluvoxamine, Furosemide.
Gabapentin, Glimepiride, Glipizide, Glyburide, Greppafloxacin.
Haloperidol, Heparin, Hydrochlorothiazide, Hydrocodone.
Ibuprofen, Imipramine, Immunosuppressants, Indinavir, Insulin, Irbesartan,
Isradipine, Isocarboxazid, Itraxonazole.
Ketoconazole.
Lansoprazole, Levonorgestrel, Letrozole, Linezolid, Lithium, Lorazepam,
Losartan, Lovastatin.
Magaldrat, Magnesium Sulfate, Medications That Slow Blood Clotting
(anticoagulant / Antiplatelet Drugs), Meperidine, Methadone, Methylene Blue,
Methylprednisolone, Metoprolol, Mexiletine, Moclobemide.
Naproxen, Narcotics, Nifedipine, Nitrates, Nizatidine, Norethindrone,
Norfloxacin.
Olanzapine, Omeprazole, Ondansetron.
Pantoprazole, Pentazocine, Pentobarbital, Phenelzine, Phenobarbital,
Phenprocoumon, Phenylpropanolamine, Phenytoin, Pioglitazone, Piroxicam,
Prednisolone, Prednisone, Primidone, Procarbazine, Propranolol,
Pseudoephedrine.
Rabeprazole, Ranitidine, Rasagiline, Riluzole, Rosiglitazone.
Safinamide, Selegiline, Sparfloxacin, Stimulants.
Tacrine, Tamoxifen, Terbinafine, Theophylline, Tolbutamide, Toremide,
Tramadol, Tranlycypromine, Trazodone, Triazolam, Trovafloxacin.
Valproic Acid, Valsartan, Verapamil.
Warfarin.
Zileuton, Zolmitriptan, Zolpidem.

If you are using any of the substances on this list, please consult with your medical practitioner before using SLAMR™.

10. Durability:

2 years, see packaging.

11. Storage conditions:

Keep it over 4 °C and under 25 °C
and away from direct sunlight.

12. Packaging (type and content):

12 grams: 1pcs | single packet of SLAMR™
1 Packet.

60 grams: 5pcs x 12 grams | box of SLAMR™
1 box, 5 Packets.

13. Side effects:

Normal:

- Increased energy.
- Mood changes.
- Sensation of being warm.
- Dry mouth.
- Restlessness.
- Heartburn.

Rare:

- Irregular heartbeat.
- Low blood pressure.
- Nosebleeds.
- Nausea.
- Diarrhea.
- Dizziness.
- Headaches.
- Stomach ache.
- Constipation.
- Excitation.
- Difficulty sleeping.
- Rapid heartbeat.
- Drowsiness.
- Changes in heart rhythm.

14. Contact:

SLAMR International OÜ
Hobujaama 10-14, Tallinn, 10151, Estonia

If you send us regular mail, please write down your e-mail address so that we can reply to your inquiry,
Or visit: www.slamrformula.com